

# Early Birds: Fossils and Feathers

A Reading A-Z Leveled Y Benchmark Book  
Word Count: 1,240



**Reading a-z**

Visit [www.readinga-z.com](http://www.readinga-z.com)  
for thousands of books and materials.

BENCHMARK • Y


# Early Birds: Fossils and Feathers



Written by Alfred J. Smuskiewicz

[www.readinga-z.com](http://www.readinga-z.com)

# Early Birds: Fossils and Feathers



Written by Alfred J. Smuskiewicz

[www.readinga-z.com](http://www.readinga-z.com)

## Photo Credits:

Front cover, page 9: © Charles R. Knight/National Geographic Images; back cover, pages 4, 8: © ArtToday; title page, pages 3, 10, 14, 15 (background): © Jupiterimages Corporation; page 5: © Jim Zuckerman/Corbis; page 7: courtesy of National Park Service; page 11: © Louie Psihoyos/Corbis; page 12: © Ken Lucas/Visuals Unlimited; page 13 (main): © Renee Jean/Daily Journal/AP Images; page 13 (inset): © Mick Ellison; page 15 (from top to bottom): © iStockphoto.com/Pavel Losevsky; page 15 (2): © iStockphoto.com/James Benet; page 15 (3, 4, 6): © iStockphoto.com; page 15 (5): © iStockphoto.com/Olga Axyutina

Early Birds: Fossils and Feathers  
Level Y Benchmark Book  
© Learning A-Z  
Written by Alfred J. Smuskiewicz


All rights reserved.

[www.readinga-z.com](http://www.readinga-z.com)

## Correlation

### LEVEL Y

Fountas & Pinnell	T
Reading Recovery	40
DRA	40



A fossil of the flightless bird Diatryma

### Table of Contents

Introduction ..... 4

Information from Fossils..... 5

Prehistoric Birds..... 8

Birds from Dinosaurs?..... 11

Conclusion ..... 14

Glossary..... 16

Index ..... 16

### Introduction

Most of the **species** of animals that have ever lived on Earth suffer extinction. When the conditions of the environment change, such as when the climate cools or the quantity of food decreases, a species may die out if it cannot **adapt** to the new conditions. **Paleontologists** have learned about a wide variety of such extinct organisms from fossils—evidence of prehistoric life preserved in rock or other material.

Some fossils are simple marks that an animal produced while moving, such as footprints or



These fossilized remains of a prehistoric bird contain many secrets of the past.

trails left in the ground. Others are hollowed-out **impressions** of an animal’s entire body made in rock. Still others are preserved remains of an animal’s body, such as bones or shells.

Fossils show how new kinds of species developed over eons. For example, scientists have discovered fossils that lead them to believe that birds developed from reptiles more than 150 million years ago; those reptiles may have been dinosaurs.

## Information from Fossils

When the vast majority of animals die, the **decomposition** activities of bacteria and fungi cause their bodies to gradually break down and disappear. However, if conditions are suitable, a dead animal may transform into a fossil, leaving behind clues about its physical characteristics and how it lived.


Typically, an animal that becomes immortalized as a fossil is buried in sediment—such as mud or sand—immediately after it dies.



John Day Fossil Beds in Oregon is typical of places fossils are found.

This is more likely to happen in or near rivers, the sea, or other bodies of water. Water that carries minerals, such as calcium, then soaks into microscopic spaces inside the bones of the body. As more and more of the bone tissue dissolves and decays away, increasing amounts of minerals take the place of the bone. In time, a rocky duplicate of the animal's skeleton is left.

When the body decays completely away, impressions of an animal's body form—showing such features as feathers or scales. All that then remains is the hollow space where the animal's body was, surrounded by sediment.



The bones and other characteristics of a fossil tell scientists what the animal looked like. These features also enable scientists to compare the animal with species living today. Similarities and differences between the features in the fossil and those of living organisms may reveal how the extinct creature behaved. For example, if the bones in the fossil are similar to bones in bird wings today, maybe the animal was capable of sustained flight. Such traits may also indicate that modern birds are related to this extinct animal.

Paleontologists use different procedures to determine how old a fossil is—that is, to learn when the animal captured as a fossil lived.

Because sediment accumulates year after year, fossils found in deeper sediment are older than fossils in sediment closer to Earth's surface. Scientists obtain their best estimates of a fossil's age by analyzing certain chemicals in the rock that contains the fossil.



Paleontologists discover fossils, study the fossils, and sometimes help display the fossils in museums.

## Prehistoric Birds

Paleontologists have unearthed numerous fossils of extinct prehistoric birds. The earliest



known bird fossils are of a primitive reptile-like bird called *Archaeopteryx* (AHR-kee-OP-tuhr-ihks), which lived approximately 150 million years ago. These fossils offer compelling evidence that birds **descended** from reptiles.

Fossils of *Archaeopteryx*, such as this one, are among the most important fossils ever discovered.

### How Old Is That Fossil?


The main chemical technique that paleontologists use to estimate the age of fossils is called *radioisotope dating*. This method is based on the fact that chemicals called radioactive isotopes break down to form other chemicals at a known rate over time. By comparing the amount of radioactive isotopes left in a fossil with the amount of their breakdown products, scientists can calculate how long this decay process has been going on. That calculation, in turn, tells scientists how long ago the fossil formed.

The fossils of *Archaeopteryx* show that this crow-sized animal had some characteristics resembling birds and others resembling reptiles. Like modern birds, *Archaeopteryx* had feathers, wings, and a “wishbone” (a forked bone in the upper chest). However, like a reptile, it had teeth and a long, bony tail. It also had three “fingers” with claws on each wing.

*Archaeopteryx* probably flew rather poorly. Scientists base that conclusion on the structure of the animal’s sternum (breastbone) seen in fossils. The sternum of *Archaeopteryx* was flat. Modern birds have a sternum with a protruding part where robust muscles used in flight are attached. Without such a sternum, *Archaeopteryx* would have lacked powerful flight muscles.



*Archaeopteryx* (top) and pterosaurs (left) were some of the first flyers besides insects.



Paleontologists have discovered several fossils younger than those of *Archaeopteryx* that depict other birds. These fossils reveal how birds developed progressively modern traits over time.

*Hesperornis* (hes-pur-OR-niss) and *Ichthyornis* (ik-thee-OR-niss) were two kinds of aquatic birds that lived approximately 90 million years ago. Fossils of *Hesperornis* show that this bird looked like a large loon, with big webbed feet to assist it in swimming. It also had a beak lined with tiny teeth to enable it to catch fish. However, *Hesperornis* had only **rudimentary** wing bones, so it could not fly. *Ichthyornis* resembled a gull, with elongated pointed wings. It was probably an excellent flyer that dived into the sea to capture fish.


Fossils prove that many of the main types of birds we know today had developed by about 35 million years ago. These included birds that bore resemblance to modern chickens, doves, ducks, parrots, penguins, owls, and songbirds.

## Birds from Dinosaurs?

Paleontologists theorize that fossils depicting birds with reptile-like characteristics are signs that birds developed from reptiles. Scientists have also uncovered fossils of dinosaurs that possessed feathers and other bird-like traits. These fossils provide evidence that dinosaurs were the reptiles from which birds developed.

Fossils indicate that **carnivorous** dinosaurs called *coelurosaurs* (SILL-ur-oh-sorz) were like birds in many ways—though scientists classify them as true dinosaurs. The fossils of the most birdlike members of the coelurosaur group are approximately 155 million to 135 million years old. They show that these animals were small for dinosaurs, most ranging in length from 2 to 10 feet (0.6–3 meters). They ran rapidly on two long, slender hind legs, which each had four, clawed toes. Their bones were hollow and lightweight. They had sizable eyes. Some even had feathers.

All of these traits are similar to those of birds. Since fossils exhibit these attributes, it's logical to conclude that coelurosaurs developed into birds.



The features seen in *Compsognathus* and other coelurosaur fossils bear several similarities to the skeletons of modern birds.

One of the coelurosaurs that paleontologists understand best from fossils is *Compsognathus* (komp-SOG-nuh-thus). It is known from two well-preserved fossils, about 145 million years old, which were discovered in Europe.

*Compsognathus* was one of the smallest dinosaurs that ever lived—some of these creatures were only the size of a chicken. *Compsognathus* had a long, thin neck and tail and long hind legs. On each of its short front legs, it had only two claws—an unusual feature for a dinosaur. Like other coelurosaurs, *Compsognathus* hunted by running after insects, lizards, and other diminutive animals and grabbing them with its razor-sharp claws and teeth.

Scientists know what this dinosaur ate because one of the fossils of *Compsognathus* includes the remains of its last meal in its stomach. The type of lizard seen in the stomach had extremely long legs, so this lizard must have been a fast runner. Thus, *Compsognathus* had to be quick to capture this prey.

*Compsognathus* was a small dinosaur that was similar to a fast-running, insect-catching bird.



### Fossil Bird in Its Egg

A 121-million-year-old fossil provides evidence that some prehistoric birds—unlike most birds today—could feed themselves immediately after hatching from their eggs. The fossil shows the outline of an egg with a baby bird still curled up inside. The unhatched bird had a complete set of feathers, strong-looking bones, and a large skull. Most birds today are weak and naked when they hatch and must be fed by their parents.

### Conclusion



Much has been learned about prehistoric animal life from fossils. Some of the most fascinating fossils ever uncovered prove that, ages ago, there were creatures that had some













characteristics of birds and other characteristics of reptiles. Fossils of these organisms lead scientists to theorize that birds developed from dinosaurs.

Dinosaurs roamed the Earth for more than 150 million years, but they were unable to adapt when environmental conditions changed about 65 million years ago. According to scientists, Earth's climate may have changed dramatically around that time, perhaps as the result of a **meteorite** impact. Although dinosaurs did not survive this change, birds did. Today, there are more than 9,700 species of birds thriving from the frigid Arctic and Antarctic to the torrid tropics.

Some scientists actually classify birds as living dinosaurs. So the next time you see a songbird singing outside your window, might you really be watching and listening to a dinosaur?





Era*	Period*	Life That Appeared
<b>Cenozoic</b> (65–present)	Quaternary (2–present)	Humans 
	Tertiary (65–2)	Horses, apes 
<b>Mesozoic</b> (248–65)	Cretaceous (145–65)	Flowering plants 
	Jurassic (213–145)	Birds, mammals 
	Triassic (248–213)	Dinosaurs 
<b>Paleozoic</b> (543–248)	Permian (286–248)	Seed plants 
	Carboniferous (360–286)	Reptiles, giant insects 
	Devonian (410–360)	Sharks, amphibians 
	Silurian (440–410)	Mosses, coral reefs 
	Ordovician (505–440)	Squid-like nautiloids 
	Cambrian (543–505)	Trilobites, mollusks, jawless fish 
	Precambrian time (4.5 billion to 543 million)	Bacteria, worms, jellyfish 

\*in millions of years

Scientists divide Earth's history into different eras and periods of time. During which era and period did birds first appear?

## Glossary

<b>adapt</b> ( <i>v.</i> )	to change to fit new conditions (p. 4)
<b>carnivorous</b> ( <i>adj.</i> )	meat-eating (p. 11)
<b>decomposition</b> ( <i>n.</i> )	the process of decaying and rotting (p. 5)
<b>descended</b> ( <i>v.</i> )	developed from; have as ancestors (p. 8)
<b>impressions</b> ( <i>n.</i> )	marks or prints made by pressure (p. 4)
<b>meteorite</b> ( <i>n.</i> )	chunk of stone or metal that falls to Earth's surface from space (p. 14)
<b>paleontologists</b> ( <i>n.</i> )	scientists who are experts in prehistoric life and fossils (p. 4)
<b>rudimentary</b> ( <i>adj.</i> )	incompletely developed (p. 10)
<b>species</b> ( <i>n.</i> )	group of organisms that share certain traits and can interbreed (p. 4)

## Index

<i>Archaeopteryx</i> , 8–10	radioisotope dating, 8
coelurosaurs, 11, 12	sternum, 9
<i>Compsognathus</i> , 12, 13	wings, 7, 9, 10
<i>Hesperornis</i> , 10	wishbone, 9
<i>Ichthyornis</i> , 10	